To discuss with your Health Care Provider



Be sure to review your plan benefits to determine your costs for these services

Annually: Discuss related screening with your doctor						
Annually						
Annually: Discuss with your doctor						
enings for At-Risk Patients						
One-time screening for ages 65 to 75 who have ever smoked or have a family history of AAA						
Women beginning at age 65 or older; and in younger women who have an increased risk						
Age 35+; Age 20-35 at risk Annually: All Men; Age 45+; Age 20-45 at risk Annually: All Women						
Age 45-75; Colonoscopy or fecal occult blood test or sigmoidoscopy						
Women should have a baseline mammogram age 35-40. Thereafter, every two years age 40-50; every year age 50+. At any age if recommended by physician based on risk factors.						
Women age 21-29 should have a Pap Test every 3 years. Women age 30-65 should have a Pap Test alone every 3 years or combined with HPV testing every 5 years. Women ages 65+ should discuss with their doctor.						
As indicated by history and/or symptoms. Discuss with your doctor behavioral risks.						
Ages 50-80; 20 pack smoker history, current smoker/quit within past 15 years						
Discuss with your doctor						
Discuss with your doctor						



Your local Blue Cross Blue Shield

Live a Healthy Lifestyle

- Get your annual wellness exam to review your overall health and keep follow-up visits with your doctor.
- Find out if you are at risk for health conditions such as diabetes, high cholesterol and high blood pressure.
- Get your vaccines, preventive screenings and labs.
- Human Papillomavirus (HPV)
 vaccine 3 dose series is
 recommended for men and
 women ages 19 through 26 years
 if not previously vaccinated prior
 to age 13.
- Talk with your doctor about the medications and over-thecounter/vitamins you are taking to reduce side effects and interactions.
- Get a Flu Vaccine every year to prevent illness and related hospitalizations.
- Get a COVID-19 vaccine to prevent severe illness and related hospitalizations.
 Immunocompromised people should consult their physician on the need for an additional mRNA vaccine dose.

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrg.gov, www.cdc.gov

To discuss with your Health Care Provider



Be sure to review your plan benefits to determine your costs for these services

Guidance	
Screen/Counseling: Depression, Obesity, Tobacco, Alcohol, Substance Abuse and Pregnancy	Every visit, or as indicated by your doctor
Fall Risk/Unintentional Injury/Domestic Violence Prevention/Seat Belt Use	Discuss exercise and home safety with your doctor
Medication List (including over-the-counter and vitamins) for potential interactions	Every visit, or as indicated by your doctor
Advance Directives/Living Will	Annually

Immunizations* (Routine Recommendations)					
Tetanus, Diphtheria, Pertussis (Td/Tdap)	Ages 19+: Tdap vaccine once, then a Td booster every 10 years				
Flu (Influenza)	Annually during flu season				
Pneumococcal PCV15, PCV20, PPSV23	Ages 19-64: if risk factors are present; Ages 65+: 1-2 doses (per CDC); Ages 50+:1 dose (Florida Blue Benefits)				
Shingles (Zoster)	Ages 50+: 2 doses Shingrix				
Haemophilus Influenzae Type b (HIB) Hepatitis A, Hepatitis B, Meningococcal	Ages 19+: if risk factors are present				
Human Papillomavirus (HPV), Measles/Mumps/Rubella (MMR), Varicella (Chickenpox) & Hepatitis C (HCV) Infection Screening	Physician recommendation based on past immunization or medical history				
COVID-19	Recommended for adults ages 19 and older within the scope of the authorization/approval for the particular vaccine.				



We're here to help: Call

Customer Service 1-800-FLA-BLUE (1-800-352-2583) TTY/TDD Call 711

Care Consultant Team 1-888-476-2227

Go to floridablue.com

Visit a Florida Blue Center

Go to **floridabluecenters.com** for locations or call 1-877-352-5830

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

^{*} Some immunizations are contraindicated for certain conditions, discuss with your doctor

To discuss with your Health Care Provider

Children & Adolescents (Birth – 18 years of age) Wellness Schedule

Routine Health Guide						
Wellness Exam and Autism/Development Behavioral Assessment	Newborn up to age 3: Frequent Wellness Check- ups; Age 3-18: Annual Wellness Check-up					
Body Mass Index (BMI): Height and Weight	Every visit, BMI beginning at age 2					
Blood Pressure	Annually, beginning at age 3					
Hearing/Dental/Vision Screenings (These services may not be covered by your medical benefits plan. Check your plan documents.)	Hearing: Newborn then annually beginning at Age 4; Dental: Regularly, beginning at age 1; Vision: Annually, beginning at age 3					
Recommended Screenings for At-Risk Patients						
Cholesterol Screening	Annually, beginning at age 2					
Lead test, TB, Sickle Cell and Blood Sugar	As indicated by history and/or symptoms					
HIV and other Sexually Transmitted Infections (STIs) Screening and Counseling	Discuss with your doctor based on behavioral risks					
Skin Cancer Screening	Discuss with your doctor					
Guidance						
Injury/Violence Prevention	Annually, more often if indicated by your doctor					
Diet/Physical Activity/Emotional Well-Being Counseling	Every visit					
Tobacco/Alcohol/Substance Abuse/Depression/Pregnancy Screening and Counseling	Every visit starting at age 11, earlier if indicated by your doctor					





Are your children up-todate with vaccinations?

Getting the recommended sequence of vaccinations is always a good idea to protect your child from illnesses from birth to 18 years of age. Most of these vaccinations require additional doses or boosters over time. As children grow up to become teenagers, they may come in contact with different diseases. Here are vaccines that can help protect your preteen or teen from these other illnesses and infections:

Tdap Vaccine

Age 11 or 12. Protects against tetanus (lock jaw), diphtheria and acellular pertussis (whooping cough). This is a booster shot of the same vaccine given during early childhood.

Meningococcal Vaccine

(MCV4) Two doses beginning at 11 or 12 years, with a booster dose at age 16. (MenB) Two doses beginning at ages 16-18. Protects against meningitis, sepsis (a blood infection) and other meningococcal diseases. Children with higher risk factors may need additional doses.

Contiunued next page

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

To discuss with your Health Care Provider

Immunizations*	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	24 months	4 - 6 years	11 - 12 years	13 - 14 years	15 years	16 - 18 years
Hepatitis A					2 dose series, 12-23 m				onths					
Hepatitis B	1st dose	2nd	dose		3rd dose									
Diphtheria, Tetanus, Pertussis (DTaP)			1st dose	2nd dose	3rd dose	4th dose		4th dose		5th dose				
Tetanus, Diphtheria, Pertussis (Tdap)											1st dose			
Haemophilus Influenzae Type b (Hib)			1st dose	2nd dose	3rd or 4th dose**									
Inactivated Poliovirus			1st dose	2nd dose	3rd dose				4th dose					
Measles, Mumps, Rubella (MMR)***						1st dose				2nd dose				
Varicela						1st dose				2nd dose				
Pneumococcal PCV13 and PPSV23			1st dose	2nd dose	3rd dose	e 4th dose								
Flu (Influenza)					6 months through 8 years 1 or 2 doses; 9 years and older 1 dose only									
Rotavirus			1st dose	2nd dose	3rd dose**									
Meningococcal											1st dose			booster
Human Papillomavirus (HPV)											3 dose series			
Covid 19					6 months and older per CDC guidelines for specific vaccine									

^{*}These are routine immunizations based upon cdc.gov recommendations. Range of recommended ages for catch-up or certain high-risk groups is at the doctor's discretion based on the member's family history and personal risk factors.





Your local Blue Cross Blue Shield

Human Papillomavirus

(HPV) Vaccine two or three doses over six months, beginning at ages 11–12. (Two doses if started before 15th birthday or three if started on or after 15). Protects boys and girls against HPV, which can lead to cancers and genital warts.

Flu Vaccine

Every year after six months of age. Protects individuals from getting the influenza virus.

COVID-19 Vaccine

The American Academy of Pediatrics (AAP) recommends the COVID-19 vaccination for all children and adolescents 6 months of age and older who do not have contraindications using a COVID-19 vaccine authorized for use for their age. Any authorized COVID-19 vaccine appropriate by age and health status can be used for COVID-19 vaccination in children and adolescents.

Keep your teens safe from preventable, painful and life-threatening diseases by staying in touch with your pediatrician's office or health clinic. Be sure to verify your benefits for preventive services.

Sources: These guidelines are recommendations from the following organizations and were not developed by Florida Blue. www.ahrq.gov, www.cdc.gov

Florida Blue is an Independent Licensee of the Blue Cross and Blue Shield Association.

^{**} Dosages determined by doctor with the type of brand vaccine used.

^{***} Those in an outbreak area, aged ≥12 months who previously received ≤ 2 doses should receive a dose of mumps-virus containing vaccine.